

Policy Brief



# Improving the Health Level of Farmers During the Use of Pesticides: Policy Brief

Fereshteh Mehri<sup>1</sup>, Akram Ranjbar<sup>1\*</sup>, Sahar Amirsadeghi<sup>1</sup>, Maryam Esfahani<sup>1</sup>, Elham Shiri<sup>2</sup>

<sup>1</sup>Nutrition Health Research Center, Hamadan University of Medical Sciences, Hamadan, Iran

<sup>2</sup>Department of Anatomical Sciences, School of Medicine, Hamadan University of Medical Sciences, Hamdan, Iran

Article history:

Received: June 1, 2022

Revised: June 6, 2022

Accepted: June 15, 2022

ePublished: June 30, 2022

\*Corresponding author:

Akram Ranjbar,

Email: akranjbar2015@gmail.com

com



## Abstract

According to the World Health Organization (WHO), the number of unintentional acute poisoning cases is between 3.5 and 5 million cases worldwide. Among these, 3 million cases were severe, leading to the death of 20000 people per year. In addition, 7.4 million years of life had lost every year due to diseases caused by agricultural toxins. Based on the previous evidence, the development of unauthorized supply stores and lack of proper monitoring of the permissible limit of poisons, as well as the sale of poisons without a prescription, the ignorance of farmers, the indiscriminate planting of products not resembling traditional agriculture, can be due to various factors related to the improper and indiscriminate use of pests. Pesticides and agricultural poisons have short- and long-term health effects. Globally, it is recommended that a policy brief be compiled for decision-making cases based on evidence. Therefore, to demonstrate evidence-based policy options, taking into account the existing obstacles and opportunities, it seems necessary to use strategies and intervention plans to reduce the consumption of pesticides while increasing the level of protection of farmers.

**Keywords:** Farmers, Pesticide, Health literacy, Policy brief

Please cite this article as follows: Mehri F, Ranjbar A, Amirsadeghi S, Esfahani M, Shiri E. Improving the health level of farmers during the use of pesticides: Policy brief. Avicenna J Pharm Res. 2022; 3(1):55-56. doi:10.34172/ajpr.2022.1080

## The Necessity of Writing a Policy Summary

The importance of environmental factors in health and well-being is becoming clearer every day (1,2). In recent decades, the presence of harmful chemicals in the environment has become a serious issue (3). To control pests and increase the production of agricultural products, pesticides have been widely used, which is the main problem in many countries (4). These poisons have been reported as the third most common cause of poisoning and the main cause of death due to poisoning in Iran (5). The average consumption of poisons per hectare is 400 g in Iran, while the average consumption of poisons in the world is 1.7 kilograms; in other words, the consumption of poisons in Iranian agricultural products is a quarter of the world average (6). The World Health Organization (WHO) estimates that there are between 3.5 and 5 million cases of acute unintentional poisoning worldwide each year. Among these, 3 million cases are severe, resulting in 20000 deaths per year (7). Further, 7.4 million years of life are annually lost due to diseases caused by agricultural toxins (8). Pesticides that are used for agricultural products may lead to poisoning, the occurrence of cancers, an increase in the incidence of cardiovascular disorders up to three times, an increase in the probability

of Parkinson's occurrence up to four times, and an increase in the incidence of hydrocephalus up to three times. In developing areas, unauthorized supply stores, a lack of proper supervision regarding the permissible limit of poisons and the sale of poisons without a prescription, ignorance of farmers, and indiscriminate planting of crops that are not typical of traditional agriculture and require high amounts of poisons can be related to various reasons (9). The incorrect and indiscriminate use of pesticides and agricultural poisons can produce short- and long-term health effects. Based on various studies, it is clear that all the management processes of poisons and pests in the country need quality improvement. It is globally recommended that a policy brief should be compiled for such decision-making cases based on the previous evidence (10). The study was developed to represent evidence-based policy options, taking into account the existing obstacles and opportunities, to decrease the consumption of pesticides while increasing the protection level of farmers so that health planners and policy-makers implement strategies and intervention plans with more confidence.

## Criticism of Current Conditions

Based on the studies conducted in this field, the general



level of the health of farmers at the time of using pesticides in Iran, as well as the skills of searching, understanding and evaluating, and applying health information and services, is low (10).

### Policy Recommendations

1. Increasing the monitoring of pesticides and their consumption limits
2. Using modern equipment to accurately identify residues in products
3. Informing farmers about how to maintain and use pesticides from certified supply centers
4. Using certified herbal supplements during using pesticides in order to reduce secondary complications.

### First Recommendation

*Planning and implementation level – Jihad Agriculture Organization:* It is recommended that the safe limit of exposure to pesticides should be determined for each product and periodically evaluated by monitoring; in addition, experts should be exposed to each product.

### Second Recommendation

*Planning and Implementation Level – Food and Drug Organization and Standards Department:* It is suggested that advanced devices such as LC-MASS/MASS and GC-MASS/MASS be used in periodical and field evaluations of the prepared samples in terms of the permissible limit and residual pesticide concentration.

### Third Recommendation

*Planning and implementation level – Jihad Agriculture and Environment Organization:* Periodic training should be provided by environmental experts and toxicology to farmers and poisoning workers to increase their awareness of health problems for consumers and their health risks.

### Fourth Recommendation

*Planning and implementation level – University of Medical Sciences:* It is recommended that clinical and laboratory experimental studies be conducted for the production of safe herbal supplement products for the use of spraying workers and farmers exposed to periodic contact with

pesticides.

### Competing Interests

The authors declare no conflict of interests.

### Funding

This study was supported by a grant from the Vice-chancellor of Research and Technology of Hamadan University of Medical Sciences for a student pharm thesis [140104282930].

### References

1. Damari B, Ahmadi Pishkuhi M, Abdollahi Z. Policy brief for reducing contaminants and residual pesticides in Iran's agricultural crops. *Community Health*. 2015;2(4):256-65. [Persian].
2. Mehri F, Ranjbar A, Shirafkan N, Soleimani Asl S, Esfahani M. The protective effect of resveratrol on diazinon-induced oxidative stress and glucose hemostasis disorder in rats' liver. *J Biochem Mol Toxicol*. 2022;36(7):e23063. doi: [10.1002/jbt.23063](https://doi.org/10.1002/jbt.23063).
3. Esfahani M, Rahbar AH, Soleimani Asl S, Mehri F. Resveratrol: a panacea compound for diazinon-induced renal toxicity. *Toxin Rev*. 2023;42(1):40-50. doi: [10.1080/15569543.2021.2008452](https://doi.org/10.1080/15569543.2021.2008452).
4. Mehri F, Goodarzi MT, Esfahani M. The possible protective effect of resveratrol on diazinon-induced oxidative stress and hepatic injury. *Avicenna J Med Biochem*. 2020;8(1):44-8. doi: [10.34172/ajmb.2020.06](https://doi.org/10.34172/ajmb.2020.06).
5. Rahbari A, Nazem H, Fazilati M, Mehri F. Protective effect of resveratrol against sub-acute diazinon-induced oxidative stress in rat kidney. *Koomesh*. 1400;23(6):794-800. [Persian].
6. Mostafalou S, Karami-Mohajeri S, Abdollahi M. Environmental and population studies concerning exposure to pesticides in Iran: a comprehensive review. *Iran Red Crescent Med J*. 2013;15(12):e13896. doi: [10.5812/ircmj.13896](https://doi.org/10.5812/ircmj.13896).
7. Salar-Amoli J, Ali-Esfahani T. Determination of hazardous substances in food basket eggs in Tehran, Iran: a preliminary study. *Vet Res Forum*. 2015;6(2):155-9.
8. Ghavami MB, Goli S, Mohammadi J, Vatandoost H. Susceptibility level of *Ornithodoros tholozani* (Acari: Argasidae) to some pesticides in north west of Iran. *Persian J Acarol*. 2015;4(1):83-94. doi: [10.22073/pja.v4i1.10194](https://doi.org/10.22073/pja.v4i1.10194).
9. Kheiripour N, Plarak A, Heshmati A, Soleimani Asl S, Mehri F, Ebadollahi-Natanzi A, et al. Evaluation of the hepatoprotective effects of curcumin and nanocurcumin against paraquat-induced liver injury in rats: modulation of oxidative stress and Nrf2 pathway. *J Biochem Mol Toxicol*. 2021;35(5):e22739. doi: [10.1002/jbt.22739](https://doi.org/10.1002/jbt.22739).
10. Rajabi F. Evidence-informed health policy making: the role of policy brief. *Int J Prev Med*. 2012;3(9):596-8.